

Care for Canthacur-PS Application to destroy lesion(s)

- 1) Leave tape on foot till next day, keep clean and dry.
- 2) If tape falls off do not worry, decrease your activity to help keep tape on.
- 3) If you have pain it usually starts 6-12 hours after application and lasts for 2-4 days.
- 4) You may pad the blistered lesion(s).
- 5) If you can take OTC Motrin or Tylenol, you may for pain.
- 6) If the blister is large and painful you may burn the tip of a needle to sterilize and try to puncture the blister to drain fluid.
- 7) Some people will soak in warm water and Epson salts, but this is not necessary.
- 8) If you have any signs of infection or any problems contact the office at 606-862-9900 in London, Irvin, or Mt. Vernon and 859-887-8026 in Nicholasville area.

Care for Canthacur-PS Application to destroy lesion(s)

- 1) Leave tape on foot till next day, keep clean and dry.
- 2) If tape falls off do not worry, decrease your activity to help keep tape on.
- 3) If you have pain it usually starts 6-12 hours after application and lasts for 2-4 days.
- 4) You may pad the blistered lesion(s).
- 5) If you can take OTC Motrin or Tylenol, you may for pain.
- 6) If the blister is large and painful you may burn the tip of a needle to sterilize and try to puncture the blister to drain fluid.
- 7) Some people will soak in warm water and Epson salts, but this is not necessary.
- 8) If you have any signs of infection or any problems contact the office at 606-862-9900 in London, Irvin, or Mt. Vernon and 859-887-8026 in Nicholasville area.

Care for Canthacur-PS Application to destroy lesion(s)

- 1) Leave tape on foot till next day, keep clean and dry.
- 2) If tape falls off do not worry, decrease your activity to help keep tape on.
- 3) If you have pain it usually starts 6-12 hours after application and lasts for 2-4 days.
- 4) You may pad the blistered lesion(s).
- 5) If you can take OTC Motrin or Tylenol, you may for pain.
- 6) If the blister is large and painful you may burn the tip of a needle to sterilize and try to puncture the blister to drain fluid.
- 7) Some people will soak in warm water and Epson salts, but this is not necessary.
- 8) If you have any signs of infection or any problems contact the office at 606-862-9900 in London, Irvin, or Mt. Vernon and 859-887-8026 in Nicholasville area.

Care for Canthacur-PS Application to destroy lesion(s)

- 1) Leave tape on foot till next day, keep clean and dry.
- 2) If tape falls off do not worry, decrease your activity to help keep tape on.
- 3) If you have pain it usually starts 6-12 hours after application and lasts for 2-4 days.
- 4) You may pad the blistered lesion(s).
- 5) If you can take OTC Motrin or Tylenol, you may for pain.
- 6) If the blister is large and painful you may burn the tip of a needle to sterilize and try to puncture the blister to drain fluid.
- 7) Some people will soak in warm water and Epson salts, but this is not necessary.
- 8) If you have any signs of infection or any problems contact the office at 606-862-9900 in London, Irvin, or Mt. Vernon and 859-887-8026 in Nicholasville area.